

BOUNDING THE HOMOLOGICAL FINITENESS LENGTH

GIOVANNI GANDINI

ABSTRACT. We give a criterion for bounding the homological finiteness length of certain $\mathbf{H}\mathfrak{F}$ -groups. This is used in two distinct contexts. Firstly, the homological finiteness length of a non-uniform lattice on a locally finite n -dimensional contractible CW-complex is less than n . In dimension two it solves a conjecture of Farb, Hruska and Thomas. As another corollary, we obtain an upper bound for the homological finiteness length of arithmetic groups over function fields. This gives an easier proof of a result of Bux and Wortman that solved a long-standing conjecture. Secondly, the criterion is applied to integer polynomial points of simple groups over number fields, obtaining bounds established in earlier works of Bux, Mohammadi and Wortman, as well as new bounds. Moreover, this verifies a conjecture of Mohammadi and Wortman.

1. INTRODUCTION

In 1984, Brown and Geoghegan showed that Thompson's group $\mathbf{F}$ is of type F_∞ giving the first example of a torsion-free group of type F_∞ of infinite cohomological dimension. In 1993 Kropholler introduced the class $\mathbf{H}\mathfrak{F}$ of *hierarchically decomposable groups* and proved that torsion-free $\mathbf{H}\mathfrak{F}$ -groups of type FP_∞ have finite cohomological dimension [15]. Hence, pathological examples like the group $\mathbf{F}$ cannot live inside $\mathbf{H}\mathfrak{F}$. The class $\mathbf{H}\mathfrak{F}$ is defined as the smallest class of groups containing the class of finite groups and which contains a group G whenever there is an admissible action of G on a finite-dimensional contractible CW-complex for which all isotropy groups already belong to $\mathbf{H}\mathfrak{F}$. Note that the class $\mathbf{H}\mathfrak{F}$ is very large and there are only a few groups which are known to lie outside $\mathbf{H}\mathfrak{F}$ [15, 1, 13].

The homological finiteness length $\phi(G)$ of a group G is a generalisation of the concepts of finite generability and finite presentability. The main result of this paper is a bound for the homological finiteness length of certain $\mathbf{H}\mathfrak{F}$ -groups:

Theorem. *Let G be an $\mathbf{H}\mathfrak{F}$ -group that acts on an n -dimensional contractible CW-complex with stabilisers of type FP_∞ . If G has no bound on the orders of its finite subgroups or it has infinite Bredon geometric dimension, then $\phi(G) < n$.*

Suppose now that X is a locally finite contractible finite-dimensional CW-complex. The topology of uniform convergence on compact subsets of X turns $\text{Aut}(X)$ into a locally compact group. Every lattice (see Section 3) on X lies in $\mathbf{H}\mathfrak{F}$ and we are able to apply the Theorem to the non-uniform ones. The S -arithmetic subgroups of non-commutative almost absolutely simple isotropic algebraic groups over global function fields are important examples of non-uniform lattices. An upper bound for

Date: June 14, 2012.

2010 Mathematics Subject Classification. Primary 20F65, 18G60.

$\phi(\Gamma)$ in this case is given in [11]. Our result provides the same bound as a corollary and we are able to prove a slightly strengthened version of a conjecture of Farb, Hruska, and Thomas [12, Conjecture 35].

Bux, Mohammadi and Wortman investigated the homological finiteness length of $\mathrm{SL}_2(\mathbb{Z}[t, t^{-1}])$, $\mathrm{SL}_n(\mathbb{Z}[t])$ and more generally integer polynomial points of simple groups over number fields [11, 9, 19]. These groups act on finite-dimensional contractible CW-complexes with arithmetic stabilisers. Note that a group lies in $\mathbf{H}\mathfrak{F}$ whenever it acts admissibly on a finite-dimensional contractible cell-complex with $\mathbf{H}\mathfrak{F}$ -stabilisers. Hence the groups considered by Bux, Mohammadi and Wortman lie in $\mathbf{H}\mathfrak{F}$ and we can apply our theorem also in this context. In particular we give the same bounds obtained in [11, 9, 19] and solve in positive a conjecture of Mohammadi and Wortman [19].

Acknowledgements. The author would like to thank Kai-Uwe Bux, Stefan Witzel and Kevin Wortman for the interesting discussions and comments without which this note would not be possible. Moreover, the author is grateful to his Ph.D. supervisor Brita E. A. Nucinkis for her encouragement and advice.

2. FINITENESS PROPERTIES

A group G is of type FP_n if the trivial $\mathbb{Z}G$ -module $\mathbb{Z}$ admits a resolution of finitely generated projective $\mathbb{Z}G$ -modules up to dimension n . If G is of type FP_n for every $n \geq 0$, then G is said to be of type FP_∞ . A group is of type F_n if it admits a $K(G, 1)$ with finite n -skeleton, and G is of type F_∞ if it is of type F_n for every $n \geq 0$. For a group, being finitely generated is equivalent to being of type FP_1 . A group is finitely presented if and only if it is of type F_2 . Bestvina and Brady show the existence of non-finitely presented groups of type FP_2 [3].

The *homological finiteness length* of G is defined as

$$\phi(G) = \sup\{m \mid G \text{ is of type } \mathrm{FP}_m\}.$$

A group acts *admissibly* on a CW-complex X if the set-wise stabiliser of each cell coincides with its point-wise stabiliser. A CW-complex X is a G -CW-complex if G acts admissibly on X .

Kropholler's class $\mathbf{H}\mathfrak{X}$ admits a hierarchy defined as follows. A group G belongs to $\mathbf{H}_1\mathfrak{X}$ if there exists a finite-dimensional contractible G -CW-complex X with cell stabilisers in $\mathfrak{X}$. The class $\mathbf{H}_\alpha\mathfrak{X}$ is defined by transfinite recursion:

- $\mathbf{H}_0\mathfrak{X} = \mathfrak{X}$;
- if α is a successor ordinal then $\mathbf{H}_\alpha\mathfrak{X} = \mathbf{H}_1(\mathbf{H}_{\alpha-1}\mathfrak{X})$;
- if α is a limit ordinal then $\mathbf{H}_\alpha\mathfrak{X} = \bigcup_{\beta < \alpha} \mathbf{H}_\beta\mathfrak{X}$.

The operator $\mathbf{H}$ is defined by “ G belongs to $\mathbf{H}\mathfrak{X}$ if and only if G belongs to $\mathbf{H}_\alpha\mathfrak{X}$ for some ordinal α ”.

The next lemma is a well-known criterion for finiteness that follows from [7, Theorem 2.2].

Lemma 2.1 ([7]). *Let G be a group that acts on an n -dimensional contractible CW-complex with stabilisers of type FP_∞ . Then, G is of type FP_∞ if and only if it is of type FP_n .*

The next result, due to Kropholler and Mislin, is the main ingredient in the proof of the Theorem.

Theorem 2.2 ([15, 16]). *Every $\mathbf{H}\mathfrak{S}$ -group of type FP_∞ has a bound on the orders of its finite subgroups and admits a finite-dimensional classifying space for proper actions.*

The Bredon geometric dimension of a group G is the minimal dimension of a classifying space for proper actions $\underline{E}G$. Note that for a torsion-free group the Bredon geometric dimension coincides with the geometric dimension.

Proof (of Theorem). By an application of Theorem 2.2 we obtain that G is not of type FP_∞ . Now we apply Lemma 2.1 to conclude that G is not of type FP_n . $\square$

3. NON-UNIFORM LATTICES ON LOCALLY FINITE CONTRACTIBLE CELL COMPLEXES

A subgroup Γ of a locally compact topological group G with left-invariant Haar measure μ is a *lattice* if:

- Γ is discrete, and
- $\mu(\Gamma \backslash G) < \infty$.

A lattice Γ is said to be uniform if $\Gamma \backslash G$ is compact. Let X be a locally finite contractible finite-dimensional CW-complex and let $\text{Aut}(X)$ be its full group of admissible automorphisms. Note that every subgroup $G \leq \text{Aut}(X)$ acts admissibly on the barycentric subdivision of X . Moreover, $\text{Aut}(X)$ is locally compact and so it is reasonable to talk about lattices on locally finite contractible finite-dimensional CW-complexes.

Lemma 3.1 ([2], Corollary 1.7). *Let X be a locally finite contractible finite dimensional CW-complex with vertex set $V(X)$. If Γ is a subgroup of $G = \text{Aut}(X)$, then:*

- Γ is discrete if and only if the stabiliser Γ_x is finite for each $x \in V(X)$;
- $\mu(\Gamma \backslash G) < \infty$ if and only if $\sum_{\sigma \in \Gamma \backslash X} \frac{1}{|\Gamma_\sigma|} < \infty$, where $\sigma = [\tilde{\sigma}]$. Moreover, the Haar measure μ can be normalised in such a way that for every discrete $\Gamma \leq G$, $\mu(\Gamma \backslash G) = \sum_{\sigma \in \Gamma \backslash X} \frac{1}{|\Gamma_\sigma|}$.

Theorem 3.2. *If Γ is a non-uniform lattice on a locally finite contractible CW-complex of dimension n , then $\phi(\Gamma) < n$.*

Proof. Let Γ be a non-uniform lattice on a locally finite contractible CW-complex X of dimension n . By Lemma 3.1 $\mu(\Gamma \backslash \text{Aut}(X)) = \sum_{\sigma \in \Gamma \backslash X} \frac{1}{|\Gamma_\sigma|}$. Since Γ is non-uniform, the set $\Gamma \backslash X$ is infinite and so for any m there is some $\sigma \in \Gamma \backslash X$ such that $\frac{1}{|\Gamma_\sigma|} < \frac{1}{m}$. Therefore, there is no bound on the orders of the stabilisers (which are finite), and so there is no bound on the orders of the finite subgroups of Γ .

Since $\Gamma \in \mathbf{H}_1\mathfrak{F}$ an application of the Theorem gives the result. Alternatively this final step can be achieved by noticing that the rational cohomological dimension of Γ is at most n and applying [15, Proposition] and Lemma 2.1 or directly by [18, Proposition 1]. $\square$

A first immediate application of Theorem 3.2 is a classical result [2, Pg. 32].

Corollary 3.3. *If X is a tree, then every non-uniform lattice in $\text{Aut}(X)$ is not finitely generated. More generally, if Γ is a non-uniform lattice on a product of n trees, then $\phi(\Gamma) < n$.*

Suppose now that X is a locally finite CAT(0) polyhedral complex [12], since every CAT(0) space is contractible [6, Corollary II. 1.5] we obtain the following:

Corollary 3.4. *If Γ is a non-uniform lattice on a locally finite CAT(0) polyhedral complex of dimension n , then $\phi(\Gamma) < n$.*

Remark 3.5. If a group G acts on a CW-complex X with finite stabilisers and X admits a compatible CAT(0)-metric, then X is a model for $\underline{EG}$ [5, Proposition 3].

Corollary 3.6. *A non-uniform lattice on a locally finite 2-dimensional contractible CW-complex cannot be finitely presented.*

The above solves a conjecture of Farb, Hruska, and Thomas [12, Conjecture 35].

As a final corollary to Theorem 3.2 we obtain the main result of [11]. We start by recalling some standard nomenclature. Let K be a global function field, and S be a finite non-empty set of pairwise inequivalent valuations on K . Let $\mathcal{O}_S \leq K$ be the ring of S -integers. Denote a reductive K -group by $\mathbf{G}$. Given a valuation v of K , K_v is the completion of K with respect to v . If L/K is a field extension, the L -rank of $\mathbf{G}$, $\text{rank}_L \mathbf{G}$ is the dimension of a maximal L -split torus of $\mathbf{G}$. The K -group $\mathbf{G}$ is L -isotropic if $\text{rank}_L \mathbf{G} \neq 0$. As in [11], to any K -group $\mathbf{G}$, there is associated a non-negative integer $k(\mathbf{G}, S) = \sum_{v \in S} \text{rank}_{K_v} \mathbf{G}$.

Corollary 3.7 (Theorem 1.2, [11]). *Let $\mathbf{H}$ be a connected non-commutative absolutely almost simple K -isotropic K -group. Then $\phi(\mathbf{H}(\mathcal{O}_S)) \leq k(\mathbf{H}, S) - 1$.*

Proof. Let $\mathbf{H}$ be a connected non-commutative absolutely almost simple K -isotropic K -group. Let H be $\prod_{v \in S} \mathbf{H}(K_v)$, there is a $k(\mathbf{H}, S)$ -dimensional Euclidean building X associated to H . X is a locally finite CAT(0) polyhedral complex. The arithmetic group $\mathbf{H}(\mathcal{O}_S)$ becomes a lattice of H via the diagonal embedding since $\mathbf{H}(\mathcal{O}_S)$ has finite covolume by [14, pg. 41]. Moreover, $\mathbf{H}$ is K -isotropic if and only if $\mathbf{H}(\mathcal{O}_S)$ is non-uniform by [14, Corollary 2.2.7]. An application of Theorem 3.2 completes the proof. $\square$

Remark 3.8. Theorem 3.2 gives the upper bound on the homological finiteness length of arithmetic groups over function fields, a historical overview can be found in [11]. In a recent remarkable paper [8] Bux, Gramlich and Witzel showed that $\phi(\mathbf{H}(\mathcal{O}_S)) = k(\mathbf{H}, S) - 1$. Calculating the homological finiteness length of non-uniform lattices on CAT(0) polyhedral complexes is an ambitious open problem. We conclude by mentioning that Thomas and Wortman exhibit examples of non-finitely generated non-uniform lattices on regular right-angled buildings [21]. This

shows that the upper bound of Theorem 3.2 is not sharp and in particular, that the Theorem of Bux, Gramlich and Witzel does not hold for all non-uniform lattices on locally finite CAT(0) polyhedral complexes.

4. ACTIONS ON NON-LOCALLY FINITE BUILDINGS

Corollary 4.1. *The following bounds hold:*

- (1) *The group $\mathrm{SL}_n(\mathbb{Z}[t])$ is not of type FP_{n-1}*
- (2) *The group $\mathrm{SL}_n(\mathbb{Z}[t, t^{-1}])$ is not of type $\mathrm{FP}_{2(n-1)}$*
- (3) *Let K be a number field and let $\mathcal{O}_K$ be its ring of integers. Let $\mathbf{H}$ be a connected, noncommutative, absolutely almost simple algebraic K -group whose K -rank equals k . Then $\mathbf{H}(\mathcal{O}_K[t])$ is not of type FP_k .*

Proof. The group $\mathrm{SL}_n(\mathbb{Z}[t])$ acts on a $(n - 1)$ -dimensional Bruhat-Tits building X . By [9, Lemma 2] the $\mathrm{SL}_n(\mathbb{Z}[t])$ stabilisers of cells in X are arithmetic groups. Arithmetic groups are of type VF by [4, Theorem 9.3]. By a construction of Serre every group of type VF has finite Bredon geometric dimension [20] and is of type F_∞ . In particular, arithmetic groups lie in $\mathbf{H}_1\mathfrak{F}$ and so $\mathrm{SL}_n(\mathbb{Z}[t]) \in \mathbf{H}_2\mathfrak{F}$. It is clear that $\mathrm{SL}_n(\mathbb{Z}[t])$ contains a free-abelian group of infinite countable rank, therefore the Bredon geometric dimension of $\mathrm{SL}_n(\mathbb{Z}[t])$ is infinite. An application of Theorem 3.2 gives part (1). It remains to prove part (2) and part (3). Arguing as in [10, Section 3] it is possible to show that the $\mathrm{SL}_n(\mathbb{Z}[t, t^{-1}])$ stabilisers of cells in the $2(n - 1)$ -dimensional Bruhat-Tits building are arithmetic groups as remarked at the end of [10, Section 4]. In the proof of [19, Lemma 2] the authors show that the $\mathbf{H}(\mathcal{O}_K[t])$ stabilisers of cells in the k -dimensional Bruhat-Tits building are arithmetic groups. In their proof there is no need to assume that the K -rank of $\mathbf{H}$ equals 2. Now proceed as in part (1) to obtain the desired bounds. $\square$

Remark 4.2. Part (1) was originally shown by Bux, Mohammadi and Wortman in [9]. Part (2) for $n = 2$ follows from a result of Krstić and McCool [17] and an alternative geometric proof is given by Bux and Wortman [10]. In the case of K -rank equals 2 part (3) was proved by Mohammadi and Wortman in [19]. In the same work they conjecture the validity of part (3) for higher ranks.

REFERENCES

1. Goulmira Arzhantseva, Martin R. Bridson, Tadeusz Januszkiewicz, Ian J. Leary, Ashot Minasyan, and Jacek Świątkowski, *Infinite groups with fixed point properties*, Geom. Topol. **13** (2009), no. 3, 1229–1263. MR 2496045 (2010b:20069)
2. Hyman Bass and Alexander Lubotzky, *Tree lattices*, Progress in Mathematics, vol. 176, Birkhäuser Boston Inc., Boston, MA, 2001, With appendices by Bass, L. Carbone, Lubotzky, G. Rosenberg and J. Tits. MR 1794898 (2001k:20056)
3. Mladen Bestvina and Noel Brady, *Morse theory and finiteness properties of groups*, Invent. Math. **129** (1997), no. 3, 445–470. MR MR1465330 (98i:20039)
4. A. Borel and J.-P. Serre, *Corners and arithmetic groups*, Comment. Math. Helv. **48** (1973), 436–491, Avec un appendice: Arrondissement des variétés à coins, par A. Douady et L. Hérault. MR 0387495 (52 #8337)
5. Noel Brady, Ian J. Leary, and Brita E. A. Nucinkis, *On algebraic and geometric dimensions for groups with torsion*, J. London Math. Soc. (2) **64** (2001), no. 2, 489–500. MR MR1853466 (2002h:57007)

6. Martin R. Bridson and André Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999. MR 1744486 (2000k:53038)
7. Kenneth S. Brown, *Finiteness properties of groups*, Proceedings of the Northwestern conference on cohomology of groups (Evanston, Ill., 1985), vol. 44, 1987, pp. 45–75. MR MR885095 (88m:20110)
8. K.-U. Bux, R. Gramlich, and S. Witzel, *Higher finiteness properties of reductive arithmetic groups in positive characteristic: the rank theorem*, ArXiv e-prints (2011).
9. Kai-Uwe Bux, Amir Mohammadi, and Kevin Wortman, $SL_n(\mathbb{Z}[t])$ is not FP_{n-1} , *Comment. Math. Helv.* **85** (2010), no. 1, 151–164. MR 2563684 (2010k:20066)
10. Kai-Uwe Bux and Kevin Wortman, *A geometric proof that $SL_2(\mathbb{Z}[t, t^{-1}])$ is not finitely presented*, *Algebr. Geom. Topol.* **6** (2006), 839–852 (electronic). MR 2240917 (2007d:20080)
11. ———, *Finiteness properties of arithmetic groups over finiteness properties of arithmetic groups over function fields*, *Inventiones Mathematicae* **167** (2007), 355–378.
12. Benson Farb, Chris Hruska, and Anne Thomas, *Problems on automorphism groups of nonpositively curved polyhedral complexes and their lattices*, *Geometry, rigidity, and group actions*, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 2011, pp. 515–560. MR 2807842
13. G. Gandini, *Cohomological invariants and the classifying space for proper actions*, To appear in *Groups, Geometry and Dynamics*.
14. G. Harder, *Minkowskische Reduktionstheorie über Funktionenkörpern*, *Invent. Math.* **7** (1969), 33–54. MR 0284441 (44 #1667)
15. Peter H. Kropholler, *On groups of type $(FP)_\infty$* , *J. Pure Appl. Algebra* **90** (1993), no. 1, 55–67. MR MR1246274 (94j:20051b)
16. Peter H. Kropholler and Guido Mislin, *Groups acting on finite-dimensional spaces with finite stabilizers*, *Comment. Math. Helv.* **73** (1998), no. 1, 122–136. MR MR1610595 (99f:20086)
17. Sava Krstić and James McCool, *The non-finite presentability of $IA(F_3)$ and $GL_2(\mathbb{Z}[t, t^{-1}])$* , *Invent. Math.* **129** (1997), no. 3, 595–606. MR 1465336 (98h:20053)
18. Ian J. Leary and Brita E. A. Nucinkis, *Bounding the orders of finite subgroups*, *Publ. Mat.* **45** (2001), no. 1, 259–264. MR MR1829588 (2002b:20074)
19. A. Mohammadi and K. Wortman, *On presentations of integer polynomial points of simple groups over number fields*, ArXiv e-prints (2011).
20. Jean-Pierre Serre, *Cohomologie des groupes discrets*, *Prospects in mathematics* (Proc. Sympos., Princeton Univ., Princeton, N.J., 1970), Princeton Univ. Press, Princeton, N.J., 1971, pp. 77–169. *Ann. of Math. Studies*, No. 70. MR MR0385006 (52 #5876)
21. Anne Thomas and Kevin Wortman, *Infinite generation of non-cocompact lattices on right-angled buildings*, *Algebr. Geom. Top.* (2011).

SCHOOL OF MATHEMATICS, UNIVERSITY OF SOUTHAMPTON, SOUTHAMPTON, SO17 1BJ UNITED KINGDOM

Current address: Rheinische Wilhelms-Universität Bonn, Mathematisches Institut, Endenicher Allee 60, 53115 Bonn, Germany

E-mail address: giovanni.gandini@hausdorff-center.uni-bonn.de